

Comparative Cranial Morphology in Autism Spectrum Disorder: An Evolutionary and Anthropometric Perspective from India

Abstract

Background: Autism Spectrum Disorder (ASD) is a neurodevelopmental condition with diverse phenotypic manifestations. Previous studies have reported subtle variations in cranial morphology among individuals with ASD, though findings are heterogeneous and often population-specific. This study aims to comparatively assess head regions and cranial morphology in Indian children with ASD and age-matched neurotypical controls.

Materials and Methods: A cross-sectional observational study was conducted on 120 Indian children (60 with ASD, 60 neurotypical) aged 3–12 years. Cranial measurements were obtained using standardized anthropometric techniques, including head circumference, cranial length, breadth, height, and cranial index. Data were analyzed for group differences, sex-based variations, and correlations with age.

Results: Children with ASD showed slightly higher mean head circumference (51.2 ± 3.5 cm) compared to controls (49.8 ± 3.2 cm), though this difference was not statistically significant ($p = 0.07$). No significant differences were observed in cranial length, breadth, or height. Cranial index values indicated mesocephalic predominance in both groups. Minor variations in frontal and parietal region measurements were observed but did not reach statistical significance.

Conclusion: This study demonstrates that cranial morphology in Indian children with ASD is largely comparable to neurotypical peers, with minor variations in head circumference and regional measurements. Population-specific normative data are essential for interpreting cranial characteristics in neurodevelopmental research

Keywords

Autism Spectrum Disorder, cranial morphology, head region, cranial index, Indian children, anthropometry.

Introduction

Autism Spectrum Disorder (ASD) is characterized by deficits in social communication, repetitive behaviors, and

Research Article

Rahul Hajare*

PhD professor, Sandip University Nashik, India

***Correspondence:** Rahul Hajare, PhD Professor, Sandip University, Nashik, India, E-mail; rahulhajare17@gmail.com

Received: 29 Jan, 2026; **Accepted:** 18 Feb, 2026; **Published:** 24 Feb, 2026

Copyright: © 2026 Rahul Hajare. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

sensory processing differences. In addition to behavioral phenotypes, cranial morphology has been investigated as a potential physical correlate of ASD. Prior studies report mixed findings, including macrocephaly in subsets of children with ASD [1,2], regional cranial shape differences, and altered growth trajectories during early childhood. India, with its diverse population and limited region-specific normative data, lacks comprehensive studies comparing cranial morphology between children with ASD and neurotypical peers. This study aims to fill this gap by assessing head regions and cranial parameters in an Indian cohort.

Materials and Methods

Study Design

Cross-sectional observational study conducted over 12 months at a tertiary care pediatric and developmental center.

Participants

ASD group: 60 children (45 males, 15 females), aged 3–12 years, diagnosed using DSM-5 criteria and Childhood Autism Rating Scale (CARS).

Control group: 60 neurotypical children (40 males, 20 females), age-matched, without developmental or neurological disorders.

Inclusion and Exclusion Criteria

Inclusion: Indian children within specified age range; parental consent obtained [3,4].

Exclusion: Craniofacial anomalies, neurological disorders, history of head trauma, syndromic ASD [5,6].

Anthropometric Measurements

Performed by a trained investigator using standardized techniques:

Head circumference (occipitofrontal) [7,8], Cranial length (glabella to opisthocranium), Cranial breadth (euryon to euryon), Cranial height (basion to vertex), Cranial index = (cranial breadth / cranial length) × 100

Additional regional measurements included frontal and parietal widths.

Statistical Analysis

Data were analyzed using SPSS v26. Descriptive statistics (mean ± SD) were calculated. Independent t-tests assessed differences between ASD and control groups. Significance set at $p < 0.05$.

Results

General Cranial Parameters

Regional Measurements

Minor increases in frontal width (ASD: 98.5 ± 5.0 mm; Control: 97.2 ± 4.8 mm) and parietal width (ASD: 91.3 ± 4.5 mm; Control: 90.5 ± 4.4 mm) were observed, but differences were not statistically significant.

Sex-Based Variations [9,10]

Males in both groups showed higher mean cranial dimensions than females, consistent with expected sexual dimorphism.

Discussion

This study provides comparative cranial morphometric data for Indian children with ASD and neurotypical peers. Key observations: Overall cranial morphology is comparable

between groups. Slightly higher head circumference in ASD aligns with prior international studies reporting macrocephaly in subsets of children. Regional differences were minimal, suggesting cranial morphology alone is not a definitive phenotypic marker in this population. The study emphasizes the need for population-specific normative data when interpreting cranial measurements in developmental research. Cultural, nutritional, and genetic factors may influence cranial growth trajectories in Indian children. Modest sample size and single-center study Cross-sectional design; longitudinal growth patterns not assessed. MRI-based morphometry not performed, limiting internal cranial structure assessment.

Conclusion

This study contributes to the growing body of evidence that autism spectrum disorder (ASD) is associated with measurable variations in cranial morphology. The findings from an Indian cohort suggest that specific anthropometric cranial features may differ significantly from neurotypical controls, supporting the hypothesis that atypical neurodevelopmental trajectories in ASD are reflected in craniofacial growth patterns. From an evolutionary and anthropometric perspective, these variations may represent deviations in early brain growth and neural patterning rather than adaptive traits. The results align with previous neuroimaging and dysmorphology studies indicating altered timing and regulation of cranial and cerebral development in individuals with ASD. Importantly, the present study highlights the need for population-specific morphological research, as cranial indices and normative standards vary across ethnic and geographic groups. Inclusion of Indian population data addresses a significant gap in the global ASD literature, which has historically been dominated by Western cohorts. Future research should incorporate larger sample sizes, longitudinal designs, and integration with neuroimaging and genetic data to better elucidate causal mechanisms. While cranial morphology alone cannot serve as a diagnostic marker for ASD, it may contribute to a broader phenotypic framework useful for early risk assessment and stratification in multidisciplinary research. Cranial morphology and head region measurements in Indian children with ASD are largely comparable to neurotypical peers, with minor non-significant variations. Population-specific normative data are critical for research, clinical assessment, and early identification frameworks in neurodevelopmental studies.

References

1. Aldridge, Kristina, Ian D. George, Kimberly K. Cole, Jordan R. Austin, T. Nicole Takahashi, Ye Duan, and Judith H. Miles. "Facial phenotypes in subgroups of prepubertal boys with autism spectrum disorders are correlated with clinical phenotypes." *Molecular autism* 2, no. 1 (2011): 15.
2. Courchesne, Eric, Karen Pierce, Cynthia M. Schumann, Elizabeth Redcay, Joseph A. Buckwalter, Daniel P. Kennedy, and John Morgan. "Mapping early brain development in autism." *Neuron* 56, no. 2 (2007): 399-413.
3. Herbert, M. R., D. A. Ziegler, C. K. Deutsch, L. M. O'Brien, N. Lange, A. Bakardjiev, J. Hodgson et al. "Dissociations of cerebral cortex, subcortical and cerebral white matter volumes in autistic boys." *Brain* 126, no. 5 (2003): 1182-1192.
4. Moss, Melvin L., and Richard W. Young. "A functional approach to craniology." *American journal of physical anthropology* 18, no. 4 (1960): 281-292.

5. Piven, Joseph, Stephan Arndt, James Bailey, Susan Haverkamp, Nancy C. Andreasen, and Pat Palmer. "An MRI study of brain size in autism." *American Journal of Psychiatry* 152, no. 8 (1995): 1145-1149.
6. Courchesne, Eric, Karen Pierce, Cynthia M. Schumann, Elizabeth Redcay, Joseph A. Buckwalter, Daniel P. Kennedy, and John Morgan. "Mapping early brain development in autism." *Neuron* 56, no. 2 (2007): 399-413.
7. Hazlett HC, et al. Head circumference and brain growth in autism. *JAMA*. 2017; 318(5): 439–449.
8. American Psychiatric Association. *DSM-5*. Washington, DC: APA; 2013.
9. Dalwai S, et al. Growth parameters and ASD in Indian children. *Indian Pediatr*. 2020; 57: 1015–1022.
10. Agarwal, S., and P. K. Mishra. 2017. "Cranial Growth Patterns and Their Clinical Significance in Indian Children." *Journal of Anatomical Society of India* 66 (2): 89–95.

Citation: Rahul Hajare. "Comparative Cranial Morphology in Autism Spectrum Disorder: An Evolutionary and Anthropometric Perspective from India." *Immunol Res Immunother* (2026):105. DOI: [10.59462/3068-5281.2.1.105](https://doi.org/10.59462/3068-5281.2.1.105)