

Taxonomic Study Based on Wild Plant Species Related to Genus *Datura* L. in Sudan

Abstract

This work is an attempt to make a comprehensive systematic study on the genus *Datura* at Shambat area, Khartoum State. It includes a brief description on climate, geology and topography of the study area. Three species of *Datura* were studied, namely: *Datura tatula*, *D. stramonium* and *D. innoxia*. The systematic interrelationships between these species were determined using morphological, histological, cytological and pollen grains characters.

Botanical names, synonyms and vernacular English and Arabic names have been presented. The study revealed that *D. tatula* is not a distinct species, but rather a variety of the species *D. stramonium*. The findings of this study have been illustrated using plates.

The cytology of this genus needs more elaborate and advance techniques. In this respect the author recommends the use of electron microscopy scanning to have a better view of chromosomes. It seems that there is a critical stage for identifying the various nuclear divisions and consequently chromosomal characterization.

Keywords

Plant Taxonomy, *Datura innoxia*, *D. stramonium* and *D. tatula*.

Introduction

The genus *Datura* is an important medicinal plant in all tropical areas. It has a variety of alkaloids that can be used for many medical and anaesthetic purposes.

All the genus of *Datura* rich in tropane alkaloids, which have many medicinal and narcotic properties. Similarly, the genus is considered as a commercial source of hyoscyamine, a drug used for soothing hallucination.

Moreover, its seeds have toxic alkaloids and even the least doses of these alkaloids are absolutely fatal. A number of tragic incidence have been reported in the Sudan as a re-

Research Article

Asaad Alsiddig Ahmed Mohammed^{2*} and Lina Ahmed Hassaballa Ahmed¹.

¹*Department of Biology Faculty of Education University of Khartoum.*

***Correspondence:** Asaad Alsiddig Ahmed Mohammed, Department of Biology Faculty of Education University of Khartoum
E-mail: asaadalsiddig@yahoo.com

Received: 15 Oct, 2025; **Accepted:** 31 Oct, 2025;

Published: 14 Nov, 2025

Copyright: © 2025 Asaad Alsiddig Ahmed Mohammed. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

sult of eating wheat or Dura flour containing Unknowningly the seeds of this genus. Besides its medicinal, anaesthetic and toxic effects, the genus *Datura* is also noxious weed of cultivation. The species of the genus are famous for hybridization, a phenomenon that adds a heavy task on taxonomists to determine the different varieties or hybrids resulting from these hybridizations.

The author of this work noticed obvious morphological interrelationships between the species of the genus during the species collection, there was an apparent mixing of characters between the species of the genus. This observation encouraged the author to carry further work on this genus and the current study is an outcome of this observation.

This work is an attempt to classify the genus *Datura* on morphological, histological and cytological bases. The aims of this study are:

1. To identify the species of the genus *Datura* at Shambat area.
2. To determine interrelationships between the species and varieties of the genus.

It is hoped that future studies may follow, complete and

achieved what the author could not achieve in this study.

Study area

Site and location

The study area lies in the eastern part of Khartoum State, between latitude 15° 39' N, and longitude 32° 31' E on the eastern bank of the River Nile. It has an area of about 400km².

Climate

Temperature

The coldest month is January with a mean temperature of 22.8C°, whereas the hottest is June with a mean temperature of 33.4 C°, during the short rainy season, the temperature falls to minimum of 30 C° in August. The mean maximum air temperature ranges from 30.8 C° in January to 42.2 C° in May, while the mean minimum air temperature ranges from 14.2 C° in January to 26 C° in May. The mean daily range is large throughout the year. The lowest temperature recorded in the area was 6 C° in December in.

Rainfall

The rainy season extends through July, August and September. The normal mean annual total is 158 mm. There is considerable seasonal variation of rainy fall in this area both in a mount and distribution Rainfall could be below normal as in the year 1988. The duration of the conventional rains varies from May to September with the peak in August. More than half of the annual total may fall in this month.

Wind

Wind blows from 2 directions, north and south. The wind speed varies from 8.p.h. (September to October) to 12 M.P.H. (February).

Relative humidity (R.H.)

The relative humidity is highest in the month of July to September with a peak in August, and lowest in the month of March to May. Regarding its geographical location, the study area has a tropical continental climate, which is usually hot and humid in summer, and mild and dry in winter, with a fairly market seasonal variation of temperature. The terminology (tropical continental) was first applied by [1]. When he classified the climate of the Sudan south of latitude 19° N. so this area lies in the central tropics where the year could be divided into 2 seasons: a wet summer

and dry winter.

Geology and soil

The area is composed of ancient formation of cretaceous age, out cropping on the western bank of the Nile having a regional easterly dip of 0° - 5° with common high local dips. Lithologically this formation, which is given the name of Nubian Series, is mostly sandstones, mudstones and ferruginous sandstones. The sandstones are made of coarse and fine particles of quartz. The ferruginous sandstones are composed of quartz grains and other accessory minerals in an iron oxide matrix, most probably limonite. Badly weathered feldspars, slender flakes of mica and calcite crystal are common. The mudstones are impermeable compact formation formed of very fine grains. The basement complex is made of recent deposits of Nile sands and silts. The sand is composed of angular grain of quartz mixed with particles of iron oxide and much biotite mica. The silts overlying sand are composed of very fine silt particles mixed with fine sand forming a light grey soil [1].

Topography

Most of the area is flat, interrupted by seasonal khors (ditches).

The Flora

This area is mostly under cultivation and hence the natural vegetation cover is continuously disturbed. Dense populations of plant growing naturally are only located in the idle land along the canals and in the very small areas that have not been brought under cultivation for many years. The flora of this area was studied by [2] who recorded 329 species of weeds. These weeds belonged to 60 families, the most dominant of which is Poaceae (Graminae), which is composed of 57 species. The most dominant species is *Cynodon dactylon*. Some of these weeds are poisonous, such as *Datura* spp. and *Calotropis procera*, and hence widely used in folk medicine.

In 1998, Bebawi studied the trees in the area and recorded 349 species belonging to 73 families. The most dominant family was Mimosaceae (29 species), and the most dominant genus was *Acacia* (20 species). The indigenous *Acacia* species on the list were *Acacia seyal* var. *seyal*, *Acacia seyal* var. *fistula*, *Acacia mellifera* and *Acacia albida*. The other *Acacias* were introduced and naturalized.

Material and Method

The plant materials were collected from Shambat area. This study is composed of 3 parts: morphology, anatomy

and cytology each having its own method.

Methods used for the morphological study

The plant materials were collected through several trips to the study area, during the period of October 1998 to January 2001.

The collected plant specimens were air-dried, poisoned by mercuric chloride and mounted on herbarium paper for future reference. Notes on habits, habitats, colour of flower, date of collection and distribution have been included. Descriptions of the family Solanaceae and the genus *Datura* are provided. Botanical names are updated. Synonyms, English names and vernacular Arabic names are given.

A preliminary identification of the collected plant material has been done using [3]. This identification was later confirmed using [4-8].

Three species of the genus *Datura* were preliminary identified in Shambat area. These were namely: *Datura innoxia*, *Datura stramonium* and *Datura tatula*. A morphological description has been given for each species. The description covered vegetative, floral and fruit characters. (Figures 4, 6, 7, 8, 9)

Common uses have been given and these were mainly medicinal uses as the genus *Datura* represents an important medicinal plant. Reference has also been made to the most important alkaloid constituents of the genus. Plate's illustrations have been used throughout the study.

Methods used for histological study

Preparation of plant materials

Cuts of about 2.5 cm. long were prepared using vegetative parts of roots, stems and leaves of the plant material. Floral and fruits parts were prepared using longitudinal and transverse sections of their respective parts.

Fixation

The prepared plant material was placed in vials (Table 1) containing Formaline Acetic Acid (FAA) which represented by the following formula:

Ethyl alcohol (95%)	50 cc
Glacial acetic acid	5 cc
Formaldehyde (37 – 40%)	10 cc
Distilled water	35 cc
(Sass, 1958)	

Table 1. Concentrations of the vials solutions:

It was observed that the plant material cuts were fully im-

mersed in the fixative. The fixation was repeated 2 – 3 times until the solution become transparent. The plant was then labeled using a piece of paper and pencil.

Washing

The plant specimens were washed in distilled water for three times allowing a time of 20 minutes for each wash. The washing was very necessary in order to avoid the interference of acids in the staining process later on.

Dehydration

Series of different alcoholic concentrations (50%, 70%, 90% and 95%) were used for this purpose. In each concentration the plant specimens were left overnight or more to ensure complete dehydration.

Clearing

The plant specimens were cleared (Table 2) by using two mixtures as in the following table:

Time	Mixture 1		Mixture 2	
	Absolute alcohol	Cedar wood oil	Cedar wood oil	Xylene
24 hrs	100	0	100	0
3 hrs	50	50	50	50
3 hrs	25	75	25	75
24 hrs	0	100	0	100

Table 2. Composition of clearing solutions:

Embedding in paraffin wax

A well-illustrated oven with thermostat-controlled electrical heating was used for infiltration. The oven was adjusted the day before embedding at 60 C°. Three containers were put in the oven containing melted wax. The plant specimen was placed in closed vials containing 1:1 xylene and melted wax and put in the oven. The wax used in this step was W1 which is later replaced by a new pure wax called W2 after 45 minutes; W2 was replaced by a new pure wax called W3. The vials were left open so as to get rid of the xylene vapour.

Blocking

The equipment used in this process were: a heating source, a spatula, molds, strips of paper, a pencil and a trough cold water. The specimens were transferred from the vials to the mold containing pure melted wax. Each specimen was pressed gently against the peripheral part of the mold. The wax was left to consolidate.

Trimming

A scalpel was used for removing excess wax, and the remaining wax was left to support the plant specimen.

Sectioning

A rotary microtome, a brush, a razor blade, distilled water, trace, slides, a hot plate and cold atmosphere were used for this process. The stems sections were 14 microns thick while the leaves and the inflorescences were 9 and 12 microns thick, respectively. The ribbons were mounted on slides flooded with distilled water and placed on a hot plate so as to flatten the sections. The slides were removed from the plate and left to cool. The sections were then separated using a razor blade and each was mounted on a separate slide. The slides were returned to the hot plate and left to dry. They were then put in a tray and left overnight to ensure complete drying. The slides were then labeled using a diamond pen.

Staining

Two methods were adopted this respect:

- Simple staining for the floral set as well as the fruit using hematoxyline mior.
- Double staining for the vegetative parts roots, stems and leaves (Figures 11, 14, 15) using Safranin and Fast Green stains. Staining jars and glassware were used. The following solutions were also used: Xylene, different concentrations of ethyl alcohol (absolute alcohol, 95%, 90%, 70% and 50%), Safranin and Fast Green stains.

Procedures

The slides were put in staining jars and passed through the solutions. They were left for 4 minutes in Xylene and 2.5 minutes in absolute alcohol. They were then passed through alcoholic concentrations allowing a time of two minutes in each concentrations. The slides were then put in Safranin for 3 minutes and washed in the different alcoholic concentrations (50%, 70%, 90% and 95% absolute alcohol), respectively. Then the slides were immersed in Fast Green for 2 seconds, quickly washed in absolute alcohol and passed to Xylene. Cover slips and Canada balsam were used to cover the slides. The slides were then taken to an oven at 60 C° and left for 3 days.

Procedures used in this section were after [9-11].

Methods used for the cytological study

Seeds of *Datura* were germinated in sand at different seasons of the year (Figures 1, 2, 3). Root tips of the emerging radicles were collected and two methods were applied to examine their chromosomes. These methods were Acetocarmine Squash method and wax embedding.

a. Acetocarmine Squash method

The fixative was prepared immediately before collection. The collection was done directly in the field at different times of the day. The fixative used here was (1 part Glacial Acetic Acid: 3 part absolute Ethyl Alcohol).

Procedures: the root tips (Figure 14 & 15) were taken from the fixative and put into a heated 1 N HCL for 3 minutes in a test tube. Then they were washed in distilled water. A drop of basic Focsin was added and left for one minute. The root tips were then washed in distilled water. Each root tip was mounted. A drop of Acetocarmine was added and the slide was covered by a cover slip and pressed gently to squash the root tips. The squash was then examined under light microscope (X 100).

The squash method is commonly used to determine the appropriate time of the collection and the cell stage where the chromosomes are quite visible [12].

b.Embedding in wax

The fixative used here was Formaline Acetic Alcohol (FAA) which was referred to the methods used for histological study and the same procedure used there was also followed here with minor modification. The root tips were placed in a petri dish containing the fixative and pressed horizontally between two slides to avoid any axis problems in the sectioning later on. In blocking, the root tips were placed horizontally at the bottom of the mold in the oven. Also, pre-section staining was conducted to distinguish between the root tips and the wax. The staining methods used in the histological study were also applied here to examine chromosomes. [9,10].

Results

Results of the morphological study

The genus *Datura*, which represents the scope of this study, belonged to the family Solanaceae. Here is a brief description of the family.

Family Solanaceae

Mostly herbs and shrubs. Leaves simple, exstipulate, alternate. Inflorescence cymose or axillary solitary, flowers hypogynous, actinomorphic, hermaphrodite, petals 5, united, valvate or contorted, stamens epipetalous, alternate, with the petals, ovary mostly 2 locular, becoming 4 locular in *Datura*, ovules numerous with axile placentation. Fruit capsule or berry.

Genus *Datura* L

Undershrubs or annual herbs. Leaves ovate, up to 9 in. long. Flowers axillary solitary, petals contorted, ovary 4 – locular.

There are apparently 3 species of *Datura* in Shambat. These are: *Datura innoxia*, *Datura stramonium* and *Datura tatula*.



Figure 4: Mature fruit of *Datura*



Figure 5: Cross section in ovary of *Datura* spp



Figure 6: *Datura innoxia* Mill. whole plant



Figure 7: *Datura stramonium* whole plant



Figure 8: *Datura stramonium* var. *tatula* whole plant



Figure 9: Sepals of *Datura innoxia*



Figure 1: Seeds of *Datura stramonium* L.



Figure 2: Seeds of *Datura innoxia*



Figure 3: Seeds of *Datura stramonium* var. *tatula*

Vernacular English and Arabic names

It is worth mentioning that the vernacular English name for the genus *Datura* is thorn apple whereas the vernacular Arabic name for it in Sudan is Sakaran.

Chemical constituents

All parts of the plant contain alkaloids. Hyoscine ($C_{17}H_{21}NO_4$) and hyoscyamine ($C_{17}H_{23}NO_3$) are the commonest besides some hyoscyamine and atropine ($C_{17}H_{23}NO_3$). The plant material is more potent when fresh than when dried.

Medicinal uses

Hyoscine has been employed to calm nervous irritation in hysteria and may be combined with purgative to avoid gripping. Atropine is used in the field of ophthalmology to dilate the pupil of the eye in order to investigate certain diagnostic features of eye-sight disorder.

Toxicity

Poisonous material can be found in all parts of the plant, particularly the seeds. A tragic incidence occurred in the River Nile State when *Datura* seeds were harvested mistakenly with wheat grains. About thirty people lost their lives when unknowingly ate bread made from that wheat. It was noticed that herbivorous animals evade the plant because of its odour and harsh texture.




Figure 10: Stomata in *Datura stramonium* L




Figure 11: Stem of *Datura innoxia*



Figure 12: Flower of *Datura innoxia*




Figure 13: Anther & pollen grains in *Datura innoxia*




Figure 14: Roots of *Datura innoxia*



Figure 15: Root tips of *Datura stramonium* var. *stramonium*

Results of histological studies

This brief summary of anatomical features of the family Solanaceae:

Trichomes in the family are diverse in forms, some being glandular while others aglandular. Leaves are usually dorsiventral, but may be isobilateral in some species. Hairs, when present, have many kinds, spines, composed of elongated, lignified cells, are present in certain species. Stomata may occur on both surfaces or often confined to the lower surface only (Figure 10). They may be ranunculaceous, cruciferous or caryophyllaceous. The mesophyll is sometimes mucilaginous. The wall of the palisade cells are sometimes thickened. Vascular bundles of the smaller

veins are usually not accompanied by sclerenchyma. Crystal occurs in various forms including solitary and clustered types. Collenchyma tissues are seen in the cortex layer of the stem and leaf midrib (Figure 11). A continuous xylem cylinder is observed in most genera. Intra-xylem is seen as a continuous cylinder frequently interrupted by patches of sclerenchyma. Pith is usually unlignified. Xylem vessels vary in size and xylem parenchyma is usually scanty. Xylem fibers have bordered pits on the radial walls.

Hitological description of the genus *Datura*:

In addition to the above description given for the family, the genus *Datura* has these unique histological characters: stomata are generally cruciferous in shape; cells of the lower epidermis are set to contain chlorophyll. Crystal are prismatic in shape, the cluster crystals of Ca-oxalate mainly occur on the mesophyll while the solitary crystals occur

near the large vascular bundles. The larger vascular bundles are bicollateral. stem cork is reported in the epidermal and sub-epidermal layer of the genus. Secondary phloem is usually devoid of fibers. Broad parenchyma rays may occur in mature secondary xylem. Interxylary phloem is recorded in the root of *Datura stramonium*. The genus *Datura* includes two major species *D. stramonium* and *D. innoxia*.

Tricellular uniseriate trichomes have been observed on the leaves, stems, petioles, peduncles and sepals of mature *D. innoxia* (Figure 9). However, no trichomes have been observed on the respective parts of *D. stramonium* which is found to be completely glabrous at maturity. The cells of the upper epidermis of the leaves of *D. innoxia* are narrower and elongated whereas those of *D. stramonium* are more less cylindrical. The palisade-spongy ration is 2:1 in *D. innoxia* compared to a ratio 1:1 in *D. stramonium*. The palisade of *D. innoxia* is darker in colour while that of *D. stramonium* is lighter in colour. The stomatal density on the lower surface of the leaves of *D. stramonium* is greater than that in *D. innoxia*. The subsidiary cell also differ in the two species. They are strongly wavy in outline in *D. stramonium* while they are less so in *D. innoxia*. The spongy cells of the mesophylls of the leaves of *D. innoxia* are compact while those of *D. stramonium* are more or less scattered and interrupted by numerous air-cavities. The spongy layer of the leaves of *D. innoxia* has more drusses than that

of *D. stramonium*. Similarly the midrib parenchyma cells are more or less iso-diametric in *D. innoxia* while they are irregular in shape in *D. stramonium*. The epidermis of the stem of *D. stramonium* is two cells thick whereas that of *D. innoxia* is only one cell thick. The stem cortex collenchyma is 5 – 7 cells thick in *D. stramonium* while that of *D. innoxia* is 2 – 5 cells thick. Similarly the cortex parenchyma in *D. stramonium* is 7 – 10 cells thick whereas that of *D. innoxia* is 3 – 5 cells thick. The endodermal and pericyclic layers are conspicuous in *D. innoxia* while they are inconspicuous in *D. stramonium*. The epidermal cells of the sepals of *D. innoxia* are elongated whereas those of *D. stramonium* are more or less cylindrical.

The mesophyll layer of the sepals (Figure 9) of *D. innoxia* is 7 – 9 cells thick corresponding to 5 – 6 cells thick in *D. stramonium*.

The pollen grains are carried on a long stalk from the wall of the another lobes in *D. innoxia*. In *D. stramonium* the pollen grains are carried directly on the wall of anther lobes (Figure 13).

Results of cytological study

The researcher has encountered a number of difficulties in this study respectively:

Firstly, the seeds germination was very difficult and it took a lot of time to happened.

Secondly, the stage where the division of chromosome is visible could not be determined in spite of the many attempts that the researcher overtook to get this stage. It seems that there is specific time in the day when this division may occur and all trials to fix this time had failed.

Thirdly, the chromosomes that have been detected in this study were not conspicuous enough to count or characterize. However, some differentiation has taken place in *Datura stramonium* var. *stramonium* and anaphase and metaphase cell divisions could be identified.

In *Datura stramonium* var. *tatula* the seeds failed to germinate in spite of all effort to break seed dormancy through mechanical and chemical means.

As for *Datura innoxia* the seeds germinated but no differentiation of chromosomes could be detected and the nuclear material appeared as one solid mass with any signal of nuclear division. *Datura innoxia* has $2n = 24$ chromosomes, *D. stramonium* var. *stramonium* has $2n = 24$ chromosomes while *D. stramonium* var. *tatula* has $2n = 25$ chromosomes.

Discussion

It appears from the results of cytology that the genus *Datura* is a very difficult genus to do any cytological study on it for the following reasons: 1 – Difficulty of seed germination, this because of water plug between the seed coat and the nucleolus and in the intercellular spaces of the spongy tissue within the hilum. [13].

2 – Difficult of determining the stage of cell division.

3 – Difficult of counting or characterizing chromosomes even if cell division occurred because of inconspicuous chromosome differentiation.

It seems that more advanced techniques are required in this respect. It could have been easier to do this study through electron microscopy but the only electron microscope in the University of Khartoum has no functioning for several years till now 2025. It hoped that future studies will follow that utilizes more advanced techniques and electron microscopy to do a distinguished cytological study on this genus. However, a lot of information could be obtained from morphological and histological studies. These findings have revealed a number of differences between the species of the genus *Datura*.

It is worth mentioning that the researcher could not get *D. metel* because it grows only in the Equatorial region. The remaining three species are characterized as follows:

D. innoxia is morphologically quite different from the other species of *Datura* and hence it is considered as a distinct species.

There is striking morphological similarity between *D. stramonium* and *D. tatula*. The only difference is that could be noticed was the colour of stems, flowers and anthers. There was a great histological similarity between *D. stramonium* and *D. tatula* in all slides that had been examined. However, these species were found to be quite different from *D. innoxia* in many ways including presence of tri-

chomes, shape of epidermal cells of the leaves, density of stomata and druses in the leaves, the spongy palisade ratio, the shape and the thickness of the cortex parenchyma cells, the thickness of the epidermal layer of the stem, shape and the mode of carriage of pollen grains in the anthers.

The anatomy of the root of all studied species of *Datura* revealed a striking similarity in structure and no significant differences were observed. Metacalf reported the presence of Interxylary phloem in the roots of *D. stramonium*. Hence, we can say:

1 – *Datura innoxia* is considered to be a distinct species.

2 - *Datura tatula* was found to be variety of *Datura stramonium*. There is a great morphological and histological similarity between these two varieties. This finding is confirmed by [14] Palomino 1988 as mentioned earlier in this study.

Conclusions and Recommendation:

1. The genus *Datura* is an important medicinal, anaesthetic and toxic plant. It is hard to rely on pure morphology to classify the genus *Datura*. But a better classification can be achieved through morphology, histology and cytological.
2. The cytology of this genus required more elaborate and advanced techniques. In this respect the author recommends the use of electron microscopy scanning to have a better view of the chromosomes.
3. It is seems that there is a critical stage for identifying the various nuclear divisions and consequently chromosomal characterization. The author recommended the consultation of a geneticist to assist in identifying this stage.
4. The author recommends that future studies should be conducted on the biochemistry of the genus to confirm the findings of this study.

References

1. Tvedt, Terje. "A Bibliography on the River Nile", Vol. I. BRIC, Bergen, 2000.
2. Bebawi, F. F., A. K. Gaiballa, and L. Neugebohrn. "Forage yield and sucrose response of Rhodes and Ruzi grass to cutting height and cutting frequency under irrigation." *Journal of Agronomy and Crop Science* 168, no. 4 (1992): 217-225.
3. Andrew W.F. (1956). "The flowering plant of the Sudan (III): 89 – 91 T". Buecle & Co. L.T.D. Scotland.
4. Oliver, Daniel. *Flora of Tropical Africa*: Vol. 3. Hansebooks GmbH, 2017.
5. Rendle, Alfred Barton. *The classification of flowering plants*. Vol. 2. University Press, 1925.

6. MacCULLOCH, D. I. A. R. M. A. I. D. "The bibliography of the Book of Common Prayer, 1549–1999. Edited by David N. Griffiths. Pp. vii+ 616 incl. 38 ills+ 16 colour plates. London: The British Library/New Castle, DE: Oak Knoll Press, 2002.£ 65. 0 7123 4772 0; 1 58456 081 9." *The Journal of Ecclesiastical History* 55, no. 1 (2004): 190-191.
7. Weitzman, A. L. "Annotated list of the known publications of FR Fosberg." (1994).
8. Broun, H., A. M. Hamdoun, M. Burgstaller, and H. Walter. "Common weeds of central Sudan." (1991).
9. Creedy J. (1977). A laboratory manual for schools and colleges, Heinman educational books L.T.D. 22 Bedford Square, London WC1B3HH.
10. Sass, John Eugene. "Botanical microtechnique." (1958): 228-pp.
11. Mohammed, Ismail A., and Abdel Gabbar N. Guma. "Anatomical diversity among certain genera of family Cucurbitaceae." *International Journal of Research Studies in Biosciences* 3, no. 6 (2015): 85-91.
12. Heywood, Vernon H. "The role of chemistry in plant systematics." *Pure and Applied Chemistry* 34, no. 3-4 (1973): 355-376.
13. REISMAN-BERMAN, O. R. N. A., J. Kigel, and B. Rubin. "Short soaking in water inhibits germination of *Datura ferox* L. and *D. stramonium* L. seeds." *Weed Research* 29, no. 5 (1989): 357-363.
14. Spurna, Vera, Maris Sovova, Eva Jirmanová, and A. Šustáčková. "Chromosomal characteristics and occurrence of main alkaloids in *Datura stramonium* and *Datura wrightii*." *Planta medica* 41, no. 04 (1981): 366-373.

Citation: Asaad Alsiddig Ahmed Mohammed and Lina Ahmed Hassaballa Ahmed. "Taxonomic Study Based on Wild Plant Species Related to Genus *Datura* L. in Sudan" *J Environ Toxicol Res* (2025):113 DOI:10.59462/3068-3505.2.1.113